

EIGHT® INSECT CONTROL GARDEN & HOME READY TO USE

- **Indoor/Outdoor Use**
- **Protects Houseplants, Roses, Flowers, Indoor Landscaping, Residential Greenhouses**
- **Kills Fungus Gnats, Scale, Whiteflies, Thrips, Aphids, Spider Mites, Mealy Bugs and More**

EPA Est. No. 4-NY-1

EPA Reg. No. 4-406

Buyers Guarantee Limited to Label Claims.

©Bonide Products, Inc. All Rights Reserved

Bonide Products, Inc.
6301 Sutliff Road
Oriskany, NY 13424

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Not for use on plants being grown for sale or for other commercial use, or for commercial seed production or for research purposes.

SHAKE WELL BEFORE USING - For use, turn nozzle to spray position, point away from face and squeeze trigger to spray.

HOUSEPLANTS

To kill insects on Indoor Landscaping direct the spray mist to hit as many insects as possible. Indoor: To Protect Plants in Residential Greenhouses and Lathouses, and for Indoor Landscaping.

Plants:	Pests:	How to Use:
Ageratum, Ardisia, Aster, Baby's Breath, Begonia, Birdsnest Fern, Bleeding Heart, Carnation, Chrysanthemum, Citrus, Coleus, Crown of Thorns, Cockscomb, Cyclamen, Dracaena, Dumbcane, English Ivy, Fuchsia, Gladiolus, Grape Ivy, Juniper, Lily, Marigold, Orchid, Palm, Pansy, Peperomia, Petunia, Philodendron, Piggy-Back Plant, Poinsettia, Portulaca, Pothos, Prayer Plant, Purple Passion, Rabbits Foot Fern,	Aphids, Armyworms, Citrus Blackfly, Corn Earworms, Exposed Thrips, Fall Cankerworms, Fungus Gnats, Inchworms, Japanese Beetles, Leafminers, Leafrollers, Mealybugs, Scales, Solder Mites, Spring Cankerworms, Whiteflies	Contact and residual: Use sufficient spray to obtain full coverage of all foliage. If application is to be made indoors, remove infested plants from living areas prior to application. Apply in an area not likely to be occupied by people or pets (particularly fish). After spraying, plants can be returned to original location. Do not use more product than recommended, as this may cause some plant injury. Use with care on ferns, palms and fuchsia. As with all plants listed above, especially carnations, chrysanthemums, and roses, there exists a wide variety of species and cultivars, and therefore, a wide

Plants Cont.:	Pests Cont.:	How to Use Cont.:
Rose, Schefflera, Snapdragon, Snake Plant, Statice, Velvet Plant, Verbena, Zinnia		variety of susceptibility to injury. To ascertain possible phytotoxicity, spray a few plants as described above, wait 24 hours, and check for possible injury.
Chrysanthemums	Armyworms, Cabbageworms, Cutworms, Leafrollers	Repeat applications may be necessary to control citrus blackflies on non-bearing citrus, Adult Japanese beetles on roses.

OUTDOOR APPLICATION

DIRECTIONS FOR APPLYING TO ROSES & FLOWERS

Roses & Flowers:	Pests:	How to Use:
Ageratum, Aster, Ardisia, Azalea, Baby's Breath, Begonia, Bird's-nest Fern, Bleeding Heart, Carnation, Chrysanthemum, Coleus, Common Ninebark and Snowberry, Crown of Thorns, Cockscomb, Cyclamen, Dracaena, Dumbcane, English Ivy, Exacum, Fuchsia, Gladiolus, Grape Ivy, Gold Bells, Hypoestes, Ivy, Juniper, Lilac, Marigold, Mock-Orange, Nannyberry, Orchid, Palm, Pansy, Pea Shrub, Peperomia, Petunia, Philodendron, Piggy-Back Plant, Poinsettia, Portulaca, Pothos, Prayer Plant,	Aphids, Armyworms, Bagworms, Cabbage Worms, Cicadas, Corn earworms, Cutworms, Exposed Thrips, Fall Cankerworms, Fall Webworms, Fungus Gnat, Inchworms, Japanese Beetles, Leafminers, Leafrollers, Mealybugs, Scale Crawlers, Spring Cankerworms, Spider Mites, Whiteflies,	Apply as a thorough spray, wetting the plants to the dripping point. Try to hit underside of leaves and penetrate dense foliage. Spray in the late afternoon or evening, when the temperature ranges from 50° to 75°F and when there is little or no wind. Spray at the first sign of insects. Repeat as necessary; use intervals of 4-8

OUTDOOR APPLICATION

Roses & Flowers Cont.:	Pests Cont.:	How to Use Cont.:
Purple Passion, Rabbits Foot Fern, Rhododendron, Rose, Schefflera, Snapdragon, Snake Plant, Statice, Velvet Plant, Verbena, Zinnia	Rose Chafer, Psyllids, Scale Insects	days. Application can be made up to the day of collection.

On Tent caterpillars, application should be made when caterpillars are young and tents are first noticed. For best results, apply in the late afternoon or evening, when caterpillars have returned to their tents. A few caterpillars may be seen escaping from the tent; however, they should die within several hours.

STORAGE AND DISPOSAL

Storage: Store and transport in an upright position. Do not contaminate water, food or feed by storage or disposal. Store in cool, dry area away from heat or open flame. **Disposal:** *If empty* - Do not reuse this container. Place in trash or offer for recycling if available. *If partly filled* - Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION - Wash after handling and before smoking or eating. Avoid contamination of feed and foodstuffs. Remove pets and birds and cover fish aquaria before spraying. Do not use on humans. Do not allow children or pets to contact treated areas until surfaces are dry.

ENVIRONMENTAL HAZARDS - This pesticide is extremely toxic to fish. Do not apply directly to water. Avoid contamination of ornamental fish ponds.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.